

Crossover from BCS to composite boson (local pair) superconductivity in quasi-2D systems

E.V. Gorbar, V.M. Loktev and S.G. Sharapov*

N.N. Bogolyubov Institute for Theoretical Physics,

252143 Kiev-143, Ukraine

Abstract

The crossover from cooperative Cooper pairing to independent bound state (composite bosons) formation and condensation in quasi-2D systems is studied. It is shown that at low carrier density the critical superconducting temperature is equal to the temperature of Bose-condensation of ideal quasi-2D Bose-gas with heavy dynamical mass, meanwhile at high densities the BCS result remains valid. The evident nonmonotoneous behaviour of the critical temperature as function of the coupling constant (the energy of the two particle bound state) is a qualitative difference of quasi-2D crossover from 3D one.

Key words: crossover, BCS, bose superconductivity, quasi-2D systems

* *Corresponding author: V.M. Loktev*

Bogolyubov Institute for Theoretical Physics

Metrologicheskaya Str. 14-b

252143, Kiev-143, Ukraine

E-mail: eppaitp@gluk.apc.org

1. The problem of crossover from the BCS theory with cooperative Cooper pairing to formation and condensation of composite bosons [1, 2] (see also the review [3]) remains of interest for a long time. After the discovery of high-temperature superconductivity (HTSC) much attention was paid to this problem [4, 5, 6]. For example, in [5], using the Bethe-Salpiter equation, it was shown that real bound states in $\mathbf{r}$ -space are formed in a 2D-superconductor as the fermion chemical potential μ takes negative values.

For pure 2D superconductors, the transition from the regime of local pairs to the Cooper pairing was studied in Refs. [4, 5] in the mean field (MF) approximation. However, it is well known that this approximation for a 2D problem is not valid for instance because of the following reasons:

i) at first, by virtue of the Coleman-Mermin-Wagner-Hohenberg theorem [7] a "super-behaviour" is not possible for pure 2D systems [8] due to the fluctuations of order parameter (especially of its phase) [9]. Of course, any real system is not 2D but at least a quasi-2D one which, to be more specific, was neglected in Refs. [4, 5], where the case of $T = 0$ was considered.

ii) at second, even for 3D systems the MF approximation in the case of local pairs superconductivity proves to be wrong [6]: in fact, instead of the true critical temperature T_c it results in the value of the temperature of local pair formation.

Taking into account the Gauss fluctuations, a study of the problem of crossover was carried out for a 3D system in Ref. [6]. However, it necessary to note that the HTSCs, study of such a crossover for which is the most actual now, are not ordinary 3D systems but quasi-2D ones because of the essential (up to five orders) anisotropy of the conductivity. Therefore, from our point of view, the generalization of results of Ref. [6] for the case of quasi-2D systems would be of significant interest.

In this communication we study a model of a quasi-2D superconductor taking into account the fluctuations of order parameter in the random phase approximation using the Coleman-

Weinberg method of effective potential. We show that in the case of high carrier density n_f when $\mu \rightarrow \epsilon_F$ (ϵ_F is the Fermi energy) the critical temperature T_c is determined by the usual BCS formula and increases with the growth of attraction between fermions. In the opposite limit – the case of local pairs – T_c is determined by the formula for the temperature of Bose-condensation of quasi-2D Bose-gas [10]. It is important that in the latter case T_c decreases with the growth of such an attraction. It is likely that the intermediate region which lies between limiting cases abovementioned is most relevant to HTSCs for which the value of T_c is probably close to its maximal one.

2. The simplest model Hamiltonian for carriers (electrons or holes) in the system reads

$$H = -\psi_\sigma^\dagger(x) \left[\frac{\nabla_\perp^2}{2m_\perp} - \frac{1}{m_z d^2} \cos(i\nabla_z d) + \mu \right] \psi_\sigma(x) - V \psi_\uparrow^\dagger(x) \psi_\downarrow^\dagger(x) \psi_\downarrow(x) \psi_\uparrow(x), \quad (1)$$

where $x \equiv t, \mathbf{r}_\perp, r_z$ (with $\mathbf{r}_\perp$ being a 2D vector); $\psi_\sigma(x)$ is a fermion field in the Shrödinger representation, σ is the spin variable; $m_\perp$ is an effective mass of carriers in planes (for example, CuO_2 ones); m_z is an effective mass in z -direction; d is the interlayer distance; V is an effective local attraction inside planes; μ and n_f are determined above; one also assumes that $\hbar = k_B = 1$. The Hamiltonian proposed proves to be very convenient for study of fluctuation stabilization by weak 3D one-particle inter-plane tunneling. We omitted the two-particle tunneling considering it smaller than the one taken into account.

It is significant that the large anisotropy of conductivity cannot be identified with the similar anisotropy of the effective masses. In particular, HTSCs with large anisotropy in z -direction do not display the metal behaviour, which shows that interplane motion of particles is incoherent. But, as will be seen, for justification of approximations used below the not very large ratio $m_z/m_\perp \geq 10^2$ is already sufficient. Namely such a value takes place, for instance, in HTSC 1-2-3.

Lagrangian density connected with (1) has the following standard form:

$$L = i\psi_\sigma^\dagger \partial_t \psi_\sigma - H, \quad \psi_\sigma \equiv \psi_\sigma(x).$$

Introducing an auxiliary Hubbard-Stratonovich field $\Phi \equiv V\psi_{\uparrow}^{\dagger}\psi_{\downarrow}^{\dagger}$ and integrating over the fermions, one obtains the following effective action

$$S_{eff}(\Phi, \Phi^*) = -i\text{TrLn}G^{-1} - \frac{1}{V} \int dt d\mathbf{r} |\Phi|^2 \quad (2)$$

with

$$G^{-1} = i\frac{\partial}{\partial t}I + \left(\frac{\nabla_{\perp}^2}{2m_{\perp}} - \frac{1}{m_z d^2} \cos(i\nabla_z d) + \mu \right) \tau_3 + \frac{\tau_1 + i\tau_2}{2}\Phi + \frac{\tau_1 - i\tau_2}{2}\Phi^*, \quad (3)$$

where I is unity matrix, τ_j are the Pauli matrices. The effective potential $U_{MF}(\Phi, \Phi^*)$ which describes the system with homogeneous order parameter in the MF approximation is determined from (2) by the following relation:

$$S_{eff}(\Phi, \Phi^*) = \int dt d\mathbf{r} [T_{kin}(\Phi, \Phi^*, \nabla\Phi, \nabla\Phi^*) - U_{MF}(\Phi, \Phi^*)],$$

where T_{kin} is a power series in derivatives¹.

Applying the usual finite-temperature formalism [11], the equation $\partial U_{MF}/\partial\Phi|_{\Phi=\Phi^*=0}=0$ result in the standard gap equation

$$\frac{1}{V} = \int \frac{d\mathbf{k}}{(2\pi)^3} \frac{1}{2\xi(\mathbf{k})} \tanh \frac{\xi(\mathbf{k})}{2T_c^{MF}}, \quad (4)$$

where $\xi(\mathbf{k}) = \varepsilon(\mathbf{k}) - \mu$ with $\varepsilon(\mathbf{k}) = \mathbf{k}_{\perp}^2/2m_{\perp} + 1/(m_z d^2) \cos(k_z d)$ (reason by which we use T_c^{MF} , rather than T_c will be clarified in what follows). We would like to emphasize that under temperatures of interest, the band width in the z -direction is $m_z^{-1}d^{-2} \ll T_c^{MF}$. As for the last inequality, it is easy to see, that at $m_z \approx 10^2 m_e$ and $d = 10\text{\AA}$ the value $\hbar^2/(m_z d^2 k_B) \sim 10\text{K}$ is really more less then the usual critical temperatures in HTSC compounds.

To study Eq. (4) in the case of the local pairs superconductivity the cutoff of BCS type can not be applied. In order to eliminate the divergences in (4) it is convenient

¹For the case of a 2D system at $T = 0$, the first terms of this series was found in Ref. [5].

to introduce (see Miyake in [1] and [4, 5]) the energy of two-particle bound state $\varepsilon_b = -2W \exp(-4\pi d/m_\perp V)$, where $W = \mathbf{k}_{\perp max}^2/2m_\perp$ is the band width in the plane ². Then, expressing the left-hand side of Eq. (4) through $|\varepsilon_b|$ and passing to the limit $W \rightarrow \infty$, one can easily regularize the gap equation. The physical meaning of this limiting procedure is based on enough natural assumption that $W \gg \epsilon_F$.

Contrary to the usual BCS approach in which it is always assumed that $\mu = \epsilon_F$, here the value of chemical potential should be consistently defined from the equation

$\partial U_{MF}/\partial \mu|_{\Phi=\Phi^*=0} = -n_f$, which leads to the second, or number, equation:

$$n_F(\mu, T_c^{MF}) \equiv \int \frac{d\mathbf{k}}{(2\pi)^3} \left[1 - \tanh \frac{\xi(\mathbf{k})}{2T_c^{MF}} \right] = n_f. \quad (5)$$

In such a case one needs to solve simultaneously the system of Eqs. (4), (5) with two unknown variables – T_c^{MF} and $\mu \equiv \mu(T_c^{MF})$, respectively (in general case of arbitrary T chemical potential $\mu \equiv \mu(T) \neq \epsilon_F$).

At high concentrations n_f , such that $\mu \gg T_c^{MF}$, equality $\mu \simeq \epsilon_F$ indeed is the solution of Eq. (5), where $\epsilon_F = \pi n_f d/m_\perp$ is the Fermi energy of free 2D fermions with $\varepsilon(\mathbf{k}) \sim \mathbf{k}^2$. Taking into account the regularization procedure, it follows from (4) that

$$T_c^{MF} \simeq \gamma \pi^{-1} \sqrt{2|\varepsilon_b| \epsilon_F} \quad (6)$$

with $\gamma \simeq 1.781$. This is just well known BCS result for a 2D metal [4, 5].

In the opposite case of small concentrations, such as $-\mu \gg T_c^{MF}$, the roles of the gap (4) and number (5) equations in some sense are reversed: the Eq. (4) determines μ , while the Eq. (5) determines T_c^{MF} . Then, from the gap equation, one can obtain that $\mu = -|\varepsilon_b|/2 - m_z^{-2} d^{-4} |\varepsilon_b|^{-1}/2$. The latter except for the second term corresponds to the result of Ref. [6]. It is directly connected with the quasi-2D character of the model and (despite this term is essentially less than the first one) is very important in the case when the fluctuations are taken into account. At $|\varepsilon_b| \gg \epsilon_F$ Eq. (4) transforms into the transcendent

² Note that in the region of parameters considered these bound states form thresholdlessly.

one: $|\varepsilon_b|/2T_c^{MF} = \ln(T_c^{MF}/\epsilon_F)$. Herefrom, we directly obtain the abovementioned result: for a finite energy ϵ_F , T_c^{MF} strongly grows as V increases. Thus, for the case of small carrier density the temperature T_c^{MF} is not connected with the critical one T_c and in fact it corresponds to the temperature of composite boson dissociation [6].

3. In order to take into account the effects of composite bosons formed in the system, we apply the Coleman-Weinberg method of the effective action [12]. In this method the effective potential $U(\Phi, \Phi^*) = U_{MF}(\Phi, \Phi^*) + U^{(1)}(\Phi, \Phi^*)$, where U_{MF} is the "tree-potential" (i.e., it is obtained in the MF approximation) and $U^{(1)}$ is one-loop correction to it. At arbitrary T 's

$$U^{(1)}(\Phi, \Phi^*) = \frac{-\imath}{2} \int \frac{d^4 q}{(2\pi)^4} \ln \left[\left(\frac{1}{V} - \frac{\imath}{(2\pi)^4} \int d^4 p \text{tr}[G(p)\tau_+ G(p+q)\tau_-] \right)^2 - \left(\frac{\imath}{(2\pi)^4} \int d^4 p \text{tr}[G(p)\tau_+ G(p+q)\tau_+] \right)^2 \right], \quad p \equiv p_0, \mathbf{p}, \quad q \equiv q_0, \mathbf{q}, \quad (7)$$

where $p_0 = \imath\pi(2j+1)T$, $q_0 = \imath 2\pi l T$ (j, l - integers), $\tau_{\pm} = (\tau_1 \pm \imath\tau_2)/2$, $G(p)$ and $G(p+q)$ are one-particle Green functions from (3) in the frequency and momentum representation. Expression (7) can be useful for the study the dependence of energy gap $|\Phi|$ on the temperature. In what follows, we restrict ourselves by the case of critical line $\Phi = \Phi^* = 0$ only and investigate the effect of the fluctuations on T_c . In this case one obtains

$$U^{(1)}(\Phi = \Phi^* = 0) = -\imath \int \frac{d^4 q}{(2\pi)^4} \ln D^{-1}\{q_0, \mathbf{q}\},$$

where

$$D^{-1}\{q_0, \mathbf{q}\} = \frac{1}{V} - \frac{1}{2} \int \frac{d\mathbf{p}}{(2\pi)^3} \frac{1}{\xi(\mathbf{p} - \mathbf{q}/2) + \xi(\mathbf{p} + \mathbf{q}/2) - q_0} \times \left[\tanh \frac{\xi(\mathbf{p} - \mathbf{q}/2)}{2T_c} + \tanh \frac{\xi(\mathbf{p} + \mathbf{q}/2)}{2T_c} \right]. \quad (8)$$

Further, one has to remove the divergences from (8) applying the same procedure of regularization as in case of the gap Eq. (4).

According to [2], we introduce the phase $\delta(\omega, \mathbf{q}) \equiv -\arg D^{-1}\{\omega + \imath 0, \mathbf{q}\}$ and take into consideration the contribution of fluctuations through the number equation (5) (see also [13])

$$n_F(\mu, T_c) + 2n_B(\mu, T_c) = n_f,$$

$$n_B(\mu, T_c) \equiv \frac{1}{2} \int \frac{d\mathbf{q}}{(2\pi)^3} \int_{-\infty}^{\infty} \frac{d\omega}{\pi} n_B(\omega) \frac{\partial \delta(\omega, \mathbf{q})}{\partial \mu}, \quad (9)$$

where $n_B(\omega) \equiv [\exp(\omega/T) - 1]^{-1}$ is the Bose distribution function. One can see from (9) that the system of fermions is divided into two coexisting and dynamically bounded systems: fermi-particles, or unbounded fermions, and local pairs, or bosons. Therewith the corresponding concentrations of fermi- and bose-particles depend on T and are known only in mean.

4. Consider the influence of $n_B(\mu, T_c)$ on the behaviour of the system as the carrier density changes. Then we should consistently solve the system of Eqs. (4) and (9). At high enough carrier density such a contribution, $n_B(\mu, T_c)$ in (9), is small, and we obtain that $T_c \simeq T_c^{MF}$. It is worth to note the following: in more consistent scheme one should take into consideration the correction $\partial U^{(1)}/\partial \Phi$ in the gap equation (4), and consequently the fluctuations would modify it. However, because of the conditions $\mu = \epsilon_F \gg T_c$ and $m_z d^2 T_c \gg 1$, one can easily be convinced that this correction changes T_c rather weakly [14]. It follows, from (6) (the T_c^{MF} coincides with the expression obtained in Ref. [5] for a pure 2D superconductor due to the band narrowness in the k_z -direction) that T_c increases with the growth of couple constant V .

At small concentrations, such that $|\mu|/T_c \gg 1$, the contribution of bosons, contrary to the previous case, is dominant and so one obtains from (9) that

$$n_B(\mu, T_c) \equiv \int \frac{d\mathbf{q}}{(2\pi)^3} n_B \left(\frac{\mathbf{q}_\perp^2}{4m_\perp} + \frac{1}{2|\varepsilon_b|(m_z d^2)^2} [1 - \cos(q_z d)] \right) \simeq \frac{n_f}{2}, \quad (10)$$

i.e. all fermions are bounded between themselves forming composite bosons. It is easy to see that the boson effective mass for its motion in the plane retains the value $2m_\perp$. As to the motion between the planes, the effective boson mass increases considerably: $2|\varepsilon_b|m_z^2 d^2 (\gg m_z)$. It is worth to stress that this increasing has the dynamical character what is simply testified by the presence of $|\varepsilon_b|$. Physically, it is ensured by the one-particle character (see Eq. (1)) of the interplane tunneling. Consequently, it is connected for a pair, with its virtual breakup, for which the energy loss is of the order $|\varepsilon_b|$.

Now, one can use the formula for Bose condensation temperature of ideal quasi-2D Bose-gas [10], it is easy to write down an equation for determining T_c , which in the case under consideration takes the simple form

$$T_c \simeq \frac{\pi n d}{2m_{\perp} \ln(2T_c |\varepsilon_b| m_z^2 d^4)} = \frac{\epsilon_F}{2 \ln(2T_c |\varepsilon_b| m_z^2 d^4)}. \quad (11)$$

The last equation describes the characteristic properties of a quasi-2D superconductor with small carrier density: *i*) at first, as the critical temperature $T_c \sim \epsilon_F$, or (see above) $T_c \sim n_f$, just as it should be in 2D case (remind that in a 3D one $T_c \sim n_f^{2/3}$ [6]), and in MF approximation 2D $T_c^{MF} \sim \sqrt{n_f} \gg T_c$ [5]; *ii*) secondly, contrary to the case of 3D superconductor where T_c does not depend on V at all [6], in a quasi-2D system T_c does depend on V , namely: T_c decreases with the growth of V . As it was stated above, the reason for this is the dynamical increasing of the composite boson mass along the third direction. Thus, the growth of $|\varepsilon_b|$ (or of V , what is the same) "makes" the system more and more two-dimensional one even for the simplest case of quasi-2D metal with a local four-fermion interaction.

It is interesting to note that decreasing T_c can also take place in the case when the local pairs are bipolarons [15]; then, the increasing of coupling with phonons, which makes the pairs more massive, also leads to T_c decreasing, (rather than increasing) as the electron-phonon coupling grows.

5. Our main results are not only expressions (6) and (11) for T_c in different limiting cases, namely, in the cases of large and small V . No less important is comparison of these expressions which show that for the given density of fermions (the given ϵ_F) there is two essential regions. If $|\varepsilon_b| \ll \epsilon_F$, then even in the case of small (by absolute value n_f) densities the BCS formula is valid and T_c grows with the increasing of $|\varepsilon_b|$ (see (6)). In the opposite case, $|\varepsilon_b| \gg \epsilon_F$, T_c decreases with the growth of $|\varepsilon_b|$. It show that in the case of quasi-2D systems (it seems that the HTSCs belong to its case) $T_c(|\varepsilon_b|)$ has a maximum, and, consequently, there is a region of optimal (with respect to $|\varepsilon_b|$) values of ϵ_F for which T_c increases. We may assume that if we take into account the two-particle tunneling in (1) (if

it does not exceed the one-particle tunneling), then we obtain only the lower boundary for T_c for large $|\varepsilon_b|$. The region of $\epsilon_F \approx |\varepsilon_b|$ needs a special study because of both presence of strongly developed fluctuations and possible distinction of properties of such a Fermi-liquid from the Fermi-liquid of Landau type. Note, in conclusion, that the fluctuations might also enhance non-monotonic character of the T_c behaviour versus n_f obtained for many-layers HTSC models [16].

Acknowledgments

We are grateful to Profs. V.P. Gusynin, E.A. Pashitskii, I.I. Ukrainskii and Dr. I.A. Shovkovy for numerous discussion and fruitful remarks. The research describe in this publication was made possible in part by Grant No K5O100.

References

- [1] D.M. Eagles, Phys. Rev. 186 (1969) 456;
A.J. Leggett, in Modern Trends in Theory of Condensed Matter, edited by A. Pekalski and R. Przystawa (Springer-Verlag, Berlin, 1980) p.13;
K. Miyake, Progr. Theor. Phys. 69 (1983) 1794.
- [2] P. Nozieres and S. Schmitt-Rink, J. Low Temp. Phys. 59 (1985) 195.
- [3] R. Mićnas, J. Ranniger and S. Robaszkiewicz, Rev. Mod. Phys. 62 (1990) 113.
- [4] M. Randeria, J. Duan and L. Shieh, Phys. Rev. Lett. 62 (1989) 981; Phys. Rev. B41 (1990) 327.
- [5] E.V. Gorbar, V.P. Gusynin and V.M. Loktev, Low Temp. Phys. 19 (1993) 832.
- [6] C.A.R. Sá de Melo, M. Randeria, and J.R. Engelbrecht, Phys. Rev. Lett. 71 (1993) 3202.
- [7] S. Coleman, Comm. Math. Phys. 31 (1973) 259;
N.D. Mermin and H. Wagner, Phys. Rev. Lett. 17 (1966) 1133;
P.C. Hohenberg, Phys. Rev. 158 (1967) 383.
- [8] In the last time, purely 2D models of superconductivity are also considered. These models do not require an order parameter and are based on Berezinskii-Kosterlitz-Thouless phase transition: V.L. Berezinskii., Zh. Eksp. Teor. Fiz. 59, (1970) 907;
J. Kosterlitz and D. Thouless, J. Phys. C6 (1973) 1181.
- [9] T.M. Rice, Phys. Rev. 140 (1965) A1889.
- [10] X.-G. Wen and R. Kan, Phys. Rev. B37 (1988) 595.

- [11] R. Schrieffer, Theory of Superconductivity (Benjamin, NY, 1964).
- [12] V.A. Miransky, Dynamical Symmetry Breaking in Quantum Field Theory (World Scientific Co., Singapore, 1993).
- [13] S. Schmitt-Rink, C.M. Varma, A.E. Ruckenstein, Phys. Rev. Lett. 63 (1989) 445.
- [14] I.E. Dzyaloshinskii and E.I. Kats, Zh. Eksp. Teor. Fiz. 55 (1968) 2373.
- [15] A.S. Alexandrov and A.B. Krebs, Usp. Fiz. Nauk, 162, No. 5 (1992) 1.
- [16] E.V. Gorbar, V.M. Loktev and S.G. Sharapov, Low Temp. Phys. 21, N.4 (1995)